



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:21 PM UTC

PDB ID : 3MVM / pdb_00003mvm
Title : P38 Alpha Map Kinase complexed with pyrrolotriazine inhibitor 7V
Authors : Sack, J.S.
Deposited on : 2010-05-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	6% 72% 16% •• 8%
1	B	366	3% 76% 12% • 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitogen-activated protein kinase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2718	1743	467	496	12			
1	B	332	Total	C	N	O	S	0	0	0
			2697	1732	462	491	12			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	expression tag	UNP Q16539
A	-4	ALA	-	expression tag	UNP Q16539
A	-3	HIS	-	expression tag	UNP Q16539
A	-2	HIS	-	expression tag	UNP Q16539
A	-1	HIS	-	expression tag	UNP Q16539
A	0	HIS	-	expression tag	UNP Q16539
A	1	HIS	-	expression tag	UNP Q16539
B	-5	MET	-	expression tag	UNP Q16539
B	-4	ALA	-	expression tag	UNP Q16539
B	-3	HIS	-	expression tag	UNP Q16539
B	-2	HIS	-	expression tag	UNP Q16539
B	-1	HIS	-	expression tag	UNP Q16539
B	0	HIS	-	expression tag	UNP Q16539
B	1	HIS	-	expression tag	UNP Q16539

- Molecule 2 is 4-{{5-(isoxazol-3-ylcarbamoyl)-2-methylphenyl}amino}-5-methyl-N-propylpyrrolidine-2,1-f[1,2,4]triazine-6-carboxamide (CCD ID: 39P) (formula: C₂₂H₂₃N₇O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			32	22	7	3		
2	B	1	Total	C	N	O	0	0
			32	22	7	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	162	Total	O	0	0
			162	162		
3	B	188	Total	O	0	0
			188	188		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 14



- Molecule 1: Mitogen-activated protein kinase 14



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.31Å 71.34Å 72.53Å 90.00° 90.47° 90.00°	Depositor
Resolution (Å)	24.48 – 2.00 24.48 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (24.48-2.00) 99.6 (24.48-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 1.99Å)	Xtriage
Refinement program	BUSTER 2.9.4	Depositor
R, R_{free}	0.197 , 0.238 0.201 , 0.248	Depositor DCC
R_{free} test set	2338 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.564	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.002 for -h,l,k 0.017 for -h,-l,-k 0.032 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5829	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 39P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	7/2780 (0.3%)	1.30	18/3772 (0.5%)
1	B	0.98	5/2758 (0.2%)	1.23	12/3741 (0.3%)
All	All	0.98	12/5538 (0.2%)	1.27	30/7513 (0.4%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	78	MET	SD-CE	-8.33	1.58	1.79
1	A	288	MET	SD-CE	-7.76	1.60	1.79
1	B	288	MET	SD-CE	-7.25	1.61	1.79
1	A	228	HIS	ND1-CE1	6.44	1.39	1.32
1	B	107	HIS	ND1-CE1	6.31	1.38	1.32
1	A	107	HIS	ND1-CE1	5.79	1.38	1.32
1	A	57	ARG	N-CA	5.66	1.50	1.46
1	B	120	GLN	CD-OE1	5.59	1.34	1.23
1	A	301	GLN	CD-OE1	5.45	1.33	1.23
1	A	325	GLN	CD-OE1	5.38	1.33	1.23
1	B	325	GLN	CD-OE1	5.23	1.33	1.23
1	A	25	GLN	CD-NE2	-5.09	1.22	1.33

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	228	HIS	CB-CG-CD2	-7.82	121.04	131.20
1	B	107	HIS	CB-CG-CD2	-7.68	121.21	131.20
1	A	228	HIS	CB-CG-CD2	-7.51	121.44	131.20
1	A	107	HIS	CB-CG-CD2	-7.51	121.44	131.20
1	A	185	THR	CA-C-N	7.44	135.75	121.54
1	A	185	THR	C-N-CA	7.44	135.75	121.54
1	A	77	HIS	CB-CG-CD2	-7.38	121.61	131.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	104	LEU	CD1-CG-CD2	-7.00	95.39	110.80
1	A	250	ILE	CA-C-N	6.18	129.18	120.28
1	A	250	ILE	C-N-CA	6.18	129.18	120.28
1	B	278	ASN	CA-CB-CG	6.13	118.73	112.60
1	A	22	GLU	CA-C-N	5.82	128.35	120.38
1	A	22	GLU	C-N-CA	5.82	128.35	120.38
1	B	269	ASN	CA-CB-CG	5.78	118.38	112.60
1	A	278	ASN	CA-CB-CG	5.76	118.36	112.60
1	A	335	ASP	CA-CB-CG	5.70	118.30	112.60
1	B	109	MET	N-CA-C	5.65	120.45	113.50
1	B	40	ALA	N-CA-C	-5.62	102.16	110.48
1	A	174	HIS	CA-C-N	5.37	131.37	121.70
1	A	174	HIS	C-N-CA	5.37	131.37	121.70
1	A	257	ASN	CA-CB-CG	5.37	117.97	112.60
1	B	43	ASP	CA-C-N	5.31	127.93	120.28
1	B	43	ASP	C-N-CA	5.31	127.93	120.28
1	A	40	ALA	N-CA-C	-5.30	101.89	110.32
1	B	275	ILE	N-CA-CB	5.29	116.58	110.49
1	B	22	GLU	CA-C-N	5.26	127.58	120.38
1	B	22	GLU	C-N-CA	5.26	127.58	120.38
1	B	324	ASP	CA-CB-CG	5.22	117.82	112.60
1	A	227	ASP	CA-CB-CG	5.20	117.80	112.60
1	A	78	MET	N-CA-C	5.07	116.86	108.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2718	0	2723	29	0
1	B	2697	0	2705	21	0
2	A	32	0	23	2	0
2	B	32	0	23	3	0
3	A	162	0	0	1	0
3	B	188	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5829	0	5474	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:LEU:HD11	1:B:232:LEU:HD11	1.64	0.80
1:B:171:LEU:HD22	2:B:361:39P:H31	1.68	0.76
1:A:78:MET:HG3	1:A:140:TYR:HE2	1.54	0.73
1:A:238:LEU:HA	1:A:268:MET:HE2	1.73	0.69
1:B:27:LEU:HD23	1:B:41:ALA:HB2	1.77	0.66
1:A:109:MET:HG3	1:A:157:ALA:HB1	1.79	0.65
1:A:27:LEU:HD23	1:A:41:ALA:HB2	1.80	0.63
1:B:171:LEU:HD13	2:B:361:39P:H30	1.82	0.61
1:A:137:GLY:O	1:A:141:ILE:HG22	1.99	0.61
1:A:78:MET:HG3	1:A:140:TYR:CE2	2.36	0.60
1:A:238:LEU:HD12	1:A:268:MET:HE3	1.82	0.60
1:A:138:LEU:HA	1:A:141:ILE:CG2	2.32	0.59
1:A:71:GLU:HB2	1:A:171:LEU:HD21	1.85	0.58
1:B:120:GLN:HG2	1:B:121:LYS:H	1.69	0.57
1:B:64:HIS:O	1:B:68:THR:HG23	2.05	0.57
1:A:174:HIS:CE1	1:A:193:ILE:HD13	2.40	0.56
1:A:23:ARG:O	1:A:44:THR:HB	2.06	0.55
1:B:23:ARG:O	1:B:44:THR:HB	2.07	0.54
1:A:210:GLY:HA2	1:A:288:MET:HE3	1.89	0.53
1:A:197:TRP:HE1	1:A:259:ILE:HD11	1.74	0.53
1:B:210:GLY:HA2	1:B:288:MET:HE3	1.91	0.52
1:B:58:PRO:CB	1:B:68:THR:HG21	2.40	0.50
1:B:108:LEU:HD23	1:B:108:LEU:C	2.36	0.50
1:A:186:ARG:HG3	1:A:194:MET:HE1	1.94	0.49
1:B:242:PRO:HB3	1:B:246:LEU:HD23	1.95	0.49
1:A:73:ARG:HE	1:A:77:HIS:CE1	2.32	0.48
1:B:253:GLU:CD	1:B:256:ARG:HH22	2.21	0.48
1:A:117:VAL:HG21	1:A:216:LEU:CD2	2.44	0.48
1:B:62:ILE:HD12	1:B:334:ILE:HG13	1.96	0.47
1:A:109:MET:HG3	1:A:157:ALA:CB	2.42	0.47
1:B:242:PRO:HB2	1:B:247:LEU:HD13	1.96	0.47
1:A:238:LEU:CD1	1:A:268:MET:HE3	2.44	0.46
1:A:185:THR:HB	1:A:186:ARG:H	1.44	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:PRO:HB3	1:A:246:LEU:HD23	1.99	0.45
1:A:173:ARG:O	1:A:175:THR:N	2.50	0.44
1:B:35:TYR:N	3:B:415:HOH:O	2.50	0.44
1:A:62:ILE:HD12	1:A:334:ILE:HG13	1.98	0.43
1:B:171:LEU:HD13	2:B:361:39P:C30	2.48	0.43
1:A:171:LEU:HG	2:A:361:39P:C31	2.48	0.43
1:A:55:LEU:HD12	1:A:102:VAL:HG13	1.99	0.43
1:B:78:MET:CE	1:B:141:ILE:HA	2.49	0.43
1:A:103:TYR:C	1:A:104:LEU:HD12	2.44	0.42
1:A:235:ILE:HD11	3:A:448:HOH:O	2.19	0.42
1:B:55:LEU:HD12	1:B:102:VAL:HG13	2.03	0.41
1:B:35:TYR:HD1	3:B:397:HOH:O	2.02	0.41
1:A:71:GLU:HG3	2:A:361:39P:H31	2.02	0.41
1:B:192:GLU:HA	1:B:197:TRP:CD1	2.56	0.41
1:A:138:LEU:HA	1:A:141:ILE:HG22	2.02	0.40
1:B:220:ARG:HE	1:B:220:ARG:HB2	1.71	0.40
1:A:253:GLU:H	1:A:253:GLU:CD	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/366 (90%)	317 (96%)	8 (2%)	4 (1%)	10	6
1	B	326/366 (89%)	319 (98%)	7 (2%)	0	100	100
All	All	655/732 (90%)	636 (97%)	15 (2%)	4 (1%)	21	17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	118	LYS
1	A	263	THR
1	A	264	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/325 (92%)	274 (91%)	26 (9%)	9	6
1	B	297/325 (91%)	275 (93%)	22 (7%)	13	9
All	All	597/650 (92%)	549 (92%)	48 (8%)	11	8

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	THR
1	A	44	THR
1	A	62	ILE
1	A	78	MET
1	A	102	VAL
1	A	108	LEU
1	A	141	ILE
1	A	151	LEU
1	A	162	CYS
1	A	173	ARG
1	A	185	THR
1	A	186	ARG
1	A	194	MET
1	A	204	VAL
1	A	232	LEU
1	A	241	THR
1	A	247	LEU
1	A	251	SER
1	A	252	SER
1	A	262	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	275	ILE
1	A	285	LEU
1	A	291	LEU
1	A	293	SER
1	A	321	ASP
1	A	333	LEU
1	B	44	THR
1	B	62	ILE
1	B	79	LYS
1	B	102	VAL
1	B	108	LEU
1	B	113	LEU
1	B	114	ASN
1	B	119	CYS
1	B	145	ASP
1	B	151	LEU
1	B	194	MET
1	B	197	TRP
1	B	199	HIS
1	B	204	VAL
1	B	220	ARG
1	B	229	ILE
1	B	232	LEU
1	B	263	THR
1	B	275	ILE
1	B	285	LEU
1	B	293	SER
1	B	333	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	HIS
1	A	310	GLN
1	B	77	HIS
1	B	100	ASN
1	B	114	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	39P	A	361	-	33,35,35	1.51	5 (15%)	37,49,49	1.81	11 (29%)
2	39P	B	361	-	33,35,35	1.52	5 (15%)	37,49,49	1.79	10 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	39P	A	361	-	-	4/20/20/20	0/4/4/4
2	39P	B	361	-	-	5/20/20/20	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	361	39P	C18-C19	3.06	1.46	1.40
2	B	361	39P	C18-C19	2.99	1.46	1.40
2	B	361	39P	C21-C22	2.77	1.43	1.39
2	A	361	39P	C21-C22	2.72	1.43	1.39
2	B	361	39P	C27-N26	-2.69	1.35	1.39
2	A	361	39P	C6-C5	2.62	1.51	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	361	39P	C6-C5	2.61	1.51	1.43
2	A	361	39P	C27-N26	-2.56	1.36	1.39
2	B	361	39P	C3-N4	2.36	1.41	1.34
2	A	361	39P	C3-N4	2.31	1.40	1.34

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	361	39P	C8-C7-C6	-3.55	104.73	106.42
2	A	361	39P	C8-C7-C6	-3.44	104.78	106.42
2	A	361	39P	C8-C9-N1	3.28	109.97	107.20
2	A	361	39P	C31-C27-N28	-3.26	106.97	112.20
2	B	361	39P	C8-C9-N1	3.24	109.94	107.20
2	B	361	39P	C31-C27-N28	-3.23	107.02	112.20
2	A	361	39P	C18-N17-C5	-3.09	118.91	128.45
2	B	361	39P	C18-N17-C5	-2.95	119.32	128.45
2	A	361	39P	O29-N28-C27	2.91	106.74	103.71
2	B	361	39P	O29-N28-C27	2.85	106.68	103.71
2	B	361	39P	C3-N2-N1	2.71	122.44	113.72
2	A	361	39P	C3-N2-N1	2.70	122.44	113.72
2	A	361	39P	C13-N12-C10	2.53	128.02	122.47
2	B	361	39P	C13-N12-C10	2.46	127.87	122.47
2	A	361	39P	N2-C3-N4	-2.30	119.24	127.47
2	B	361	39P	N2-C3-N4	-2.29	119.25	127.47
2	B	361	39P	C24-N26-C27	-2.18	122.71	126.41
2	A	361	39P	C19-C18-N17	2.14	122.24	118.82
2	A	361	39P	O25-C24-C22	-2.07	116.80	120.90
2	B	361	39P	C19-C18-N17	2.02	122.05	118.82
2	A	361	39P	C23-C18-N17	-2.00	116.71	121.95

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	361	39P	N28-C27-N26-C24
2	A	361	39P	C31-C27-N26-C24
2	B	361	39P	N28-C27-N26-C24
2	B	361	39P	C31-C27-N26-C24
2	B	361	39P	N12-C13-C14-C15
2	A	361	39P	C23-C18-N17-C5
2	B	361	39P	C23-C18-N17-C5
2	B	361	39P	C19-C18-N17-C5

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	361	39P	C19-C18-N17-C5


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	361	39P	2	0
2	B	361	39P	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/366 (91%)	0.43	22 (6%) 24 23	19, 34, 61, 72	0
1	B	332/366 (90%)	0.08	11 (3%) 49 48	15, 30, 55, 77	0
All	All	667/732 (91%)	0.26	33 (4%) 35 34	15, 32, 59, 77	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	199	HIS	5.3
1	A	175	THR	5.0
1	A	30	VAL	4.6
1	B	35	TYR	4.3
1	A	263	THR	3.5
1	A	77	HIS	3.3
1	A	321	ASP	3.2
1	B	36	GLY	3.2
1	A	116	ILE	3.0
1	A	141	ILE	3.0
1	A	319	VAL	2.9
1	B	293	SER	2.8
1	A	275	ILE	2.8
1	A	117	VAL	2.7
1	A	197	TRP	2.7
1	A	122	LEU	2.7
1	A	271	ALA	2.6
1	B	30	VAL	2.6
1	B	351	PRO	2.6
1	B	198	MET	2.5
1	A	272	ASN	2.5
1	A	230	ASP	2.4
1	A	253	GLU	2.4
1	A	162	CYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	172	ALA	2.3
1	B	110	GLY	2.3
1	A	226	THR	2.2
1	A	320	ALA	2.2
1	A	119	CYS	2.2
1	B	331	ASP	2.2
1	B	197	TRP	2.1
1	A	123	THR	2.1
1	A	311	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	39P	B	361	32/32	0.95	0.07	15,25,39,43	0
2	39P	A	361	32/32	0.96	0.07	19,27,44,45	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 39P B 361:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 39P A 361:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.